

# Table of recommended dosages


## General explanations

<b>**</b>	Wherever these asterisks appear – it is recommended that the protocols are used in conjunction with each other
<b>Grey Shaded Area</b>	All Protocols that are suggested as a combined treatment are shaded in grey, and we recommend that they be used in conjunction with each other
<b>Orally</b>	Always mix the recommended drops with an equal number of activator together in a glass, wait the prescribed 3 minutes, add a mixer as per Chapter 7, and drink immediately.
<b>Transdermally</b>	Always mix the recommended drops with an equal number of activator together in a glass, wait the prescribed 3 minutes, add the prescribed amount of DMSO, wait an additional 3 minutes – then apply to the skin using your bare hands. NEVER use latex gloves. In the event of risk of contamination to the person applying the DMSO – (in the case of a patient with open sores from AIDS/HIV – apply to an area that does not present open sores, or use a wooden spatula or the back of a solid silver / stainless steel spoon to apply.
<b>Intravenously</b>	For intravenous protocols contact Dr McDonald 0027 82 339 9799
<b>Anally</b>	Always mix the recommended drops with an equal number of drops of ACTIVATOR – wait the prescribed 3 minutes then mix ACTIVATED solution with 30cc of water. Using a syringe and soft tube – insert into the anus. Alternatively use an enema pump.
<b>Allopathic Medicines</b>	It is quite safe to combine FAITH Drops with other medicines EXCEPT other alternative medicines. We prefer that you use ONLY one protocol at a time.
<b>DMSO</b>	Dimethyl Sulfoxide. DO NOT use in conjunction with cortisone medication. Wait 8 hours before starting DMSO protocol.
<b>Atropine</b>	DO NOT use in conjunction with DMSO protocol
<b>Adrenaline</b>	DO NOT use in conjunction with DMSO protocol

KEY:

FD = FAITH DROPS

ACT = ACTIVATOR



**ALWAYS MIXED IN A 1: 1 RATIO**

**\*\* INDICATES THAT PROTOCOLS SHOULD BE COMBINED**

<b>ILLNESS</b>	<b>FD</b>	<b>ACT</b>	<b>SCHEDULE</b>	<b>COMMENTS</b>
Abscesses / boils - Orally	3	3	Day 1 to 7 – 4 x Daily thereafter	Mix activated drops with approx. 250ml (250cc) of water and drink.
Abscessed teeth – Orally	6	6	2 x Daily – until clear	Mix activated drops with approximately 75ml (75cc) water then rinse out the mouth. Hold solution in the mouth for 60 seconds. FAITH™ Drops will not penetrate tooth enamel to relieve infection inside a tooth unless there is a crack or a hole for the drops to enter.
Aids – Orally Transdermal Intravenously Anally	-	-	-	Follow Protocols - Specialised Treatments – HIV/AIDS.
Allergies - Orally	3	3	2 x Daily – until clear	Mix activated drops with 250ml (250cc) of water and drink.
A.L.S. – Lou Gehrig Disease - Orally	3	3	3 x Daily - until tests clear	Follow Assail Protocols under Protocols and Treatments
Anaemia – Orally	3	3	Day 1 to 7 4 x Daily - thereafter 3 x Daily - until normalised	Mix activated drops with 250ml (250cc) of water and drink.
Arteriosclerosis – Orally	3	3	3 x Daily Until tests clear	Follow Assail Protocols under Protocols and Treatments
Arthritis - Orally	3	3	Day 1 to 7 4 x Daily thereafter 2 x Daily - until clear	Faith™ Drops will not repair damage done to tissue caused by arthritis. Ascertain the type of arthritis being treated as FAITH will assist in the treatment of rheumatoid arthritis, Lyme arthritis and arthritis. Mix activated drops with 250ml (250cc) of water and drink
Arthritis - Lyme - Orally	3	3	2 x Daily – until clear	Mix activated drops with 250ml (250cc) of water and drink. This is a long term treatment
Asthma - Orally	3	3	2 x Daily – until clear	Mix activated drops with 250ml (250cc) of water and drink. This is a long term treatment
Athlete's feet - Topically	10	10	4 x Daily – until clear	Mix activated drops with 250ml (250cc) of water. Apply the solution to affected area. Wash off after 3 minutes and dry well.
Bacteraemia (Blood-poisoning) - Orally	3	3	4 x Daily – until clear.	Mix activated drops with 250ml (250cc) of water and drink.
Bedsores Diabetes sores / ulcers - topically	10	10	Every 2 hours until healing begins	Mix activated drops with 60ml (60cc) of water and pour into a spray bottle. Spray bedsore with a fresh solution every 2 hours until healing takes place.
Bronchitis - Orally	3	3	4 x Daily - until clear.	Mix activated drops with 250ml (250cc) of water and drink.

<b>ILLNESS</b>	<b>FD</b>	<b>ACT</b>	<b>SCHEDULE</b>	<b>COMMENTS</b>
Bites - animal, insect, human, snake or spider - Orally	3	3	Hourly for 4 hours thereafter 2 hourly until clear	Mix activated drops with 250ml (250cc) of water and drink.
Cancer - Orally Transdermal Anally Intravenously **	-	-	-	Follow Faith OCC Protocols. Intravenous protocol available to medical practitioners only.
Candida – Thrush - Orally	3	3	Day 1 - Hourly thereafter 4 x Daily - until clear.	Mix activated drops with 175ml (175cc) water and rinse out the mouth. Do not leave this solution in the mouth for longer than 60 seconds
Candida – Thrush – Topically	10	10	3 x Daily - until clear.	Mix the activated drops with 20ml (20cc) of water. Apply the solution to affected area.
Cellulites - Orally	3	3	4 x Daily - until clear.	Mix activated drops with 250ml (250cc) of water and drink.
Cholesterol - Orally	3	3	2 x Daily – until clear.	Mix activated drops with 250ml (250cc) of water and drink.
Chicken-pox – Orally	2	2	Day 1 – Hourly thereafter 4 x Daily - until clear	Mix activated drops with 250ml (250cc) of water and drink.
Cholera - Orally	3	3	Day 1 – Hourly Day 2 - Every 2 hours not more than 4 x Daily Day 3 - Every 4 hours - not more than 2 x Daily	Mix activated drops with 250ml (250cc) of water and drink.
Common Cold - Orally	3	3	4 x Daily until clear.	Mix activated drops with 250ml (250cc) of water and drink.
Crabs – lice – Topically	10	10	Repeat every 2 hours until clear.	Dilute activated drops into 125ml (125cc) of water. Apply to area and rinse off after 3 minutes
Croup – Orally	2	2	4 x Daily until clear.	Mix activated drops with 250ml (250cc) of water and drink.
Cystitis – Orally	3	3	Day 1 - 4 x Daily thereafter 3 x Daily until clear.	Mix activated drops with 250ml (250cc) of water and drink.
Cysts – Orally	3	3	Day 1 - 4 x Daily thereafter 2 x Daily until clear	Mix activated drops with 250ml (250cc) of water and drink.
Dandruff – Topically	10	10	2 x Daily until clear.	Dilute activated drops into 175ml (175cc) of water. Apply to area and rinse off after 3 minutes.

<b>ILLNESS</b>	<b>FD</b>	<b>ACT</b>	<b>SCHEDULE</b>	<b>COMMENTS</b>
Dermatitis - Orally	3	3	4 x Daily until clear	Mix activated drops with 250ml (250cc) of water and drink.
Diabetes - Orally	-	-	-	Follow the Diabetes 1 + 2 protocols
Diarrhoea – Orally	3	3	Day 1 – Hourly thereafter Day 2 – Every 2 hours not more than 4 x Daily Day 3 – Every 4 hours not more than 2 x Daily	Mix activated drops with 250ml (250cc) of water and drink.
Diverticulitis - Orally	3	3	Day 1 - 4 x Daily thereafter 3 x Daily - until clear.	Mix activated drops with 250ml (250cc) of water and drink.
Eczema - Orally	3	3	3 x Daily until clear	Mix activated drops with 250ml (250cc) of water and drink.
Encephalitis - Orally	3	3	Day 1 - hourly thereafter 4 x Daily until clear.	Mix activated drops with 250ml (250cc) of water and drink.
Eye Infections - Stye Conjunctivitis - Iritis -  Orally **	3	3	Day 1 - hourly thereafter Day 2 - Every 2 hours Day 3 – Every 2 hours Thereafter take Maintenance Dose of 3 drops 3 x Daily for ten days	Combine with Eye Rinse Protocol as below

ILLNESS	FD	ACT	SCHEDULE	COMMENTS
Eye Infections - Stye Conjunctivitis Iritis Extreme Eye Infections  Eye Rinse Protocol **	1	1	Repeat every 2 hours until infection has cleared.	Combine with Oral Treatment as above. Follow instruction carefully as one can cause damage to the eyes. <ul style="list-style-type: none"> <li>o Add one drop of FAITH to 15ml (15cc) (1 tablespoon) of distilled water or eye drops.</li> <li>o Wash eye with this solution, apply directly into eye using an eye rinse bowl.</li> <li>o Allow drops to remain in the eye for 15 seconds. <b>set a timer. Do not guess at the time.</b></li> <li>o Then use distilled water. or eye drops to rinse the eye, ensuring all SCD has been rinsed from the area.</li> </ul> Repeat every two hours until infection has cleared. <ul style="list-style-type: none"> <li>o If you are unable or unwilling to administer the drops directly into the eyes, oral dosage will be sufficient provided the intake is increased to 3 drops hourly.</li> </ul>
Flu – Orally	3	3	Day 1 - 4 x Daily thereafter 3 x Daily until clear.	Mix activated drops with 250ml (250cc) of water and drink.
Food poisoning – Orally	3	3	Hourly for 4 hours thereafter Every two hours until clear	Mix activated drops with 250ml (250cc) of water and drink.
Gastritis - Orally	2	2	Day 1 - 4 x Daily thereafter 3 x Daily until clear	Mix activated drops with 250ml (250cc) of water and drink.
Glandular fever – Orally	3	3	Day 1 - 4 x Daily thereafter 3 x Daily until clear.	Mix activated drops with 250ml (250cc) of water and drink.
Gout – Orally	3	3	Day 1 to Day 7 – 4 x Daily thereafter 2 x Daily until clear	Mix activated drops with 250ml (250cc) of water and drink.

ILLNESS	FD	ACT	SCHEDULE	COMMENTS
Head Sores – Topically	10	10	1 x daily - repeat until clear	Mix activated drops with 250ml (250cc) of water and drink. When combed into the hair with enough solution to penetrate to the scalp it will cure most head sores in a few hours. If you use it daily it makes for a healthy scalp, but it also opens the cuticle scale of the hair and strips the natural melanin out of the hair. So, if you are not a natural blonde we suggest you use it once or twice a month. o When used daily your natural colour will be stripped from the hair and your hair will be bleached blonde.
Heart - Pericarditis Orally	1	1	Day 1 – Hourly thereafter 2 drops every 2 hours until clear.	Mix activated drops with 250ml (250cc) of water and Drink.
Hepatitis A, B & C – Orally & Intravenous	-	-	-	Follow Faith Hepatitis Protocols. Intravenous protocols available to Medical practitioners only
Herpes – Orally	3	3	4 x Daily	Mix activated drops with 250ml (250cc) of water and <b>Drink. With severe cases take the formula every two hours for several days. In extreme cases, one may have to use the drops intravenously, by calculating 1 drop per 10kg of body weight. Add UNACTIVATED drops to 500ml drip twice daily.</b>
Infected Gums – Orally	6	6	2 x Daily	Mix activated drops with 175ml (175cc) of water and use this solution to rinse the mouth. Do not leave this solution in the mouth for longer than 60 seconds.
Insect Bites – Topically	10	10	As required	Apply an activated solution topically directly onto the bites either with clean fingers or an earbud. You do not need to wash this solution off, for once you have added the activator to the drops you have negated its ability to burn your skin. When using the solution on larger areas of the body other than one or two small insect bite add about 75-125ml (75-125cc) of water to activated drops.
Jock Itch – Topically	6	6	3 x Daily – until clear.	Mix activated drops with 125ml (125cc) of water. Apply the solution with your finger tips to affected area. Rinse the area after 3 minutes as the area is very sensitive.
Kidney- Pylonephritis - Orally	2	2	4 x Daily – until clear	Mix activated drops with 250ml (250cc) of water and Drink. Confirm if Assail Protocol required.

<b>ILLNESS</b>	<b>FD</b>	<b>ACT</b>	<b>SCHEDULE</b>	<b>COMMENTS</b>
Laryngitis – Orally	2	2	4 x Daily – until clear	Mix activated drops with 250ml (250cc) of water and Drink.
Leukemia – Orally	-	-	-	Follow Faith OCC Protocols
Lupus – Orally	3	3	2 x Daily - until clear	Mix activated drops with 250ml (250cc) of water and Drink.
Lyme Disease – Orally	3	3	Day 1 to 14 Every 2 hours thereafter 3 x Daily - until clear.	Mix activated drops with 250ml (250cc) of water and Drink. This is a long term treatment. Information received to date from users of SCD on Lyme Disease show improvement in symptoms.
Malaria - Orally	-	-	-	Follow Faith Malaria Protocols
Measles	1	1	Day 1 - 2 x hourly thereafter Every 4 hours thereafter until clear.	Mix activated drops with 250ml (250cc) of water and Drink.
Melanomas – Orally Tranndermally **	-	-	-	Follow Faith OCC Protocols
Meningitis – Orally	2	2	Day 1 – Hourly thereafter 2 hourly thereafter until clear.	Mix activated drops with 250ml (250cc) of water and Drink. Confirm if Aggressive Protocol required.
Mumps – Orally	2	2	Every 3 hours until clear.	Mix activated drops with 250ml (250cc) of water and Drink.
Obesity – Orally	2	2	3 x Daily	Calculate 1 drop for every 10kg of body weight. Increase dose by 1 drop daily until full dose reached according to weight, but not exceeding a maximum of 10 drops per dose unless recommended by your Healthcare Practitioner. Mix activated drops with 250ml (250cc) of water and Drink.
Osteomyilitis – Orally	3	3	Day 1 – 2 hourly thereafter 4 hourly until clear	Mix activated drops with 250ml (250cc) of water and Drink.

<b>ILLNESS</b>	<b>FD</b>	<b>ACT</b>	<b>SCHEDULE</b>	<b>COMMENTS</b>
Otitis – Orally	2	2	4 x Daily until clear	Mix activated drops with 250ml (250cc) of water and Drink.
Pancreatitis – Orally	2	2	Day 1 to 2 4 hourly thereafter 4 x Daily until clear.	Mix activated drops with 250ml (250cc) of water and Drink.
Pneumonia – Orally	3	3	Day 1 – 4 hourly Thereafter 4 x Daily until clear.	Mix activated drops with 250ml (250cc) of water and Drink.
Pyorrhoea – Orally	6	6	2 x Daily until clear.	Activate drops and use solution to brush the teeth. Make and use a new solution each time. Do not leave this solution in the mouth for longer than 60 seconds.
Ringworm – Orally Topically **	3	3	4 x Daily until clear.	Follow Standard Dose in Protocols. Utilise Topical Application protocols
Scabies – Orally	3	3	2 x Daily until clear	Mix activated drops with 250ml (250cc) of water and Drink.
Scabies – Topically	10	10	Repeat every 3 hours until clear	Dilute with 125ml of water. Apply to area then rinse off after 3 minutes.
Septicaemia – Orally	3	3	Every hour for 1 day thereafter every 4 hours thereafter until clear.	Mix activated drops with 250ml (250cc) of water and Drink.
Shingles / Herpes - Orally	2	2	Every 2 hours for 1 day thereafter 3 drops 4 x Daily until clear.	Mix activated drops with 250ml (250cc) of water and Drink.
Sinusitis - Orally	3	3	Every 4 hours until clear.	Mix activated drops with 250ml (250cc) of water and Drink.
Spider Bites - Brown Recluse Spider – Orally **	3	3	3 x Daily until clear	Mix activated drops with 250ml (250cc) of water and Drink. Use in conjunction with Topical Protocol below.

<b>ILLNESS</b>	<b>FD</b>	<b>ACT</b>	<b>SCHEDULE</b>	<b>COMMENTS</b>
Spider Bites - Brown Recluse Spider – Topically **	10	10	3 x Daily for two days	Mix activated drops in a tot measure and wait prescribed 3 minutes. Mix in 5 drops (using an eye dropper) of DMSO into the activated solution. Wait an additional 3 minutes. Mix this solution with 5ml (5cc) of water. Invert tot measure over the spider bite and hold firmly in place for 3 minutes. This will ensure that the bite is totally submerged in the FAITH solution. In conjunction with this apply any nappy rash 5 minutes after the application of the DMSO protocol. Nappy rash ointment containing at least (higher if possible) 40% zinc oxide will kill most of the poison and reduces the pain and itching of a brown recluse spider bite.
Spider Bites - Black Widow – Orally Topically INtravenously **	3	3	3 x Daily for 2 to 3 days to aid the body in recovering	Use Standard Oral Protocol in conjunction with Topical & Intravenous Protocols. (For Medical Practitioners only)
Tuberculosis - Orally Intravenously **				Follow Faith TB Protocols
Tumors – Orally Transdermally **	-	-	-	Follow Faith OCC Protocols
Urinary tract infections Orally	3	3	4 x Daily until clear.	Mix activated drops with 175ml (175cc) of water and drink.
Ulcers – Orally	3	3	4 x Daily until clear.	Mix activated drops with 175ml (175cc) of water and drink.
Vaginal infections – Orally	3	3	4 x Daily until clear.	Mix activated drops with 175ml (175cc) of water and drink.
Vaginitis – Orally	3	3	4 x Daily until clear	Follow Faith Treatment Protocols – for Douches
Vaginosis – Orally	3	3	4 x Daily until clear	Mix activated drops with 175ml (175cc) of water and drink.
Warts – Topically	10	10	Repeat daily until wart falls off	Mix the drops and apply directly onto wart. Rinse off after 3 minutes.
Weight Control	3	3	3 x Daily until normalised	Mix activated drops with 175ml (175cc) of water and drink.
Whooping Cough – Orally	2	2	4 x Daily until clear	Mix activated drops with 175ml (175cc) of water and drink.
Yeast infection – orally	-	-	-	Follow Faith Candida Protocols